



RESEARCH ARTICLE

THREE-DIMENSIONAL EVALUATION OF THE FIRST CERVICAL VERTEBRAL MORPHOLOGY IN SKELETAL CLASS I AND III MALOCCLUSIONS IN YEMENI PATIENTS

Etehad Al-Ghola¹, Ghamdan Al-Harazi²

Orthodontics, Pedodontics and Prevention Department Faculty of Dentistry, Sana'a University, Yemen.

Article Info:

**Article History:**

Received: 11 December 2024

Reviewed: 6 January 2025

Accepted: 21 February 2025

Published: 15 March 2025

Cite this article:

Al-Ghola E, Al-Harazi G. Three-dimensional evaluation of the first cervical vertebral morphology in skeletal class i and iii malocclusions in Yemeni patients. Universal Journal of Pharmaceutical Research 2025; 10(1): 20-24.

<http://doi.org/10.22270/ujpr.v10i1.1269>

***Address for Correspondence:**

Etehad Al-Ghola, Orthodontics, Pedodontics and Prevention Department Faculty of Dentistry, Sana'a University, Yemen. Tel: +967-1-239643; E-mail: al-ghola_ete@yemen.net.ye

Abstract

Introduction: The shape of the cervical vertebrae is known to be of great importance in the field of orthodontics, as it can be used to assess skeletal maturity. This study aimed to evaluate the shape of the cervical vertebrae in individuals with class I and III skeletal malocclusion.

Methods: The research data were collected from the orthodontic patient records by analyzing cone beam computed tomography of the selected individuals before treatment (n=52) this study was conducted between January 2022 and January 2023. The individuals were divided into two groups based on the ANB angle: class I group (n=26, male=12, female=14) and class III group (n=26, male=9, female=17). The ages ranged from 18 to 30 years, and they were all Yemeni.

Results: The shape of the cervical vertebrae differs among individuals with different anteroposterior skeletal patterns. Cone beam radiographs of 52 patients were evaluated to assess the morphology of the first cervical vertebra in both Class I and Class III. Eight linear variables and one angular variable were compared in both groups, and there were significant differences between HOTDC1, LOAPC1, dorsal arch, HOTDC1, H1APC1, FOTDC1, and superior surface among individuals with different anteroposterior skeletal patterns.

Conclusion: The morphology of the cervical vertebrae was found to be influenced by the anteroposterior relationship of the maxilla to the mandible.

Keywords: Cervical vertebrae, Cone-beam computed tomography (CBCT), orthodontics, Yemen.

INTRODUCTION

The assessment of the patient's skeletal age is very important to determine the appropriate treatment plan for orthodontic patients¹. There are seven vertebrae in the cervical spine, which support the head. In addition, research has been conducted on the relationship between C1 dimensions and skull and neck position. The length of the mandible and neck position has shown a correlation; longer mandibles were associated with more cervical columns tilted toward the true horizontal². Class I, II, and III skeletal malocclusions can be identified by the anteroposterior relationship of the base³. Class III malocclusions may be manifested by skeletal and dental features, including mandibular skeletal protrusion, maxillary skeletal recession, or a combination of the two. Alveolar protrusions such as retroclined mandibular incisors and proclined maxillary incisors are common dental features⁴. Orthodontics is one of the medical specialties that has widely used 3D technologies. The use of these technologies has changed the way orthodontists and maxillofacial

surgeons diagnose, plan treatment, monitor condition, and evaluate outcomes. These technologies accurately visualize 3D anatomy by reproducing anatomical structures^{5,6}. Several biological indicators, such as cervical vertebral morphology and hand and wrist X-rays, can be used to determine a patient's growth stage^{7,8}. For facial and skull imaging, cone-beam computed tomography (CBCT) is superior to medical CT due to its variable field of view (FOV), fast scanning speed, very high resolution of patient radiation exposure, isotropic units, rapid image analysis, and image enhancement⁹. With the development of cone-beam computed tomography (CBCT) technology, many authors have used it to evaluate the morphology of the cervical vertebrae because it gives more accurate three-dimensional images and shows any abnormalities in the cervical vertebrae¹⁰.

Recently, there has been a revolution in Yemen to know many dental standards, including "the reliability of modern dental age estimation methods using X-rays among Yemeni children¹¹, dental caries and treatment

needs among children with physical disabilities¹², the effect of dental implants on increasing the colonization rate of aerobic bacteria in the oral cavity¹³, the effect of dental implants on the colonization of aerobic bacteria in the oral cavity, the antibiotic profile of common isolated aerobic bacteria¹⁴, temporomandibular dystonia: prevalence, clinical and demographic data¹⁵, and results of treatment strategies for hundreds of patients, radiographic evaluation of prominent fillings using cone beam computed tomography¹⁶, factor resolution, and the pattern of permanent tooth extraction¹⁷, the prevalence of signs of temporomandibular joint disorders in healthy edentulous individuals¹⁸, and the validity of Tanaka and Johnston's mixed dentition among Yemeni adults¹⁹. However, a comprehensive evaluation of the shape of the cervical vertebrae in cases of Class I and III malocclusion in the Yemeni population has not been performed. Therefore, the aim of this study was to investigate the shape of the cervical vertebrae in cases of Class I and III malocclusion using CT imaging technique.

SUBJECTS AND METHODS

Sample size: The study by Watanabe *et al.* 2 reported a dorsal arch measurement of 8.3 ± 1.1 mm in skeletal class II against 9.9 ± 2.6 mm in skeletal class III. The sample size was calculated using G*power 3.0.10 software with an alpha value of 0.05 and a power of 80%. A minimum sample size of 26 participants was determined via power analysis. Between January 2022 and January 2023, the sample was taken from the orthodontic patients' records using their pre-treatment CBCT (N=52). Subjects of this study were selected from the different Skeletal Classes (Class I and Class III). Patients were considered Skeletal Class I pattern when ($ANB = 3 \pm 1^\circ$), Skeletal Class III pattern when ($ANB < 1^\circ$)¹¹. Two gender-based subgroups were created from each study group: Class I group (n=26, males=12, females=14). Group Class III (n=26; 9 males, 17 females). Cone beam CT scans are used for orthodontic diagnosis and treatment planning, and the records used in this study were gathered from Sana'a University's Faculty of Dentistry. All of the volunteers were Yemeni and ranged in age from 18 to 30. There were no congenital maxillofacial abnormalities among the individuals.

Reliability: For accuracy, a well-trained assessor was used on the E-A scanner (Ez3Dplus2009; Ewoosoft, Co., Ltd., Hwaseong, Korea) and the analysis and reading were performed twice within a two-week period to eliminate measurement errors.

Measurements: This study measures eight linear measurements and one angular measurement to assess the morphology of cervical vertebrae with different sagittal skeletal patterns. Those measurements were as follows:

1. Horizontal outer anteroposterior (AP) diameter of the first cervical vertebra (C1) (mm) (HOAPCI) (Figure 1).
2. Horizontal inner anteroposterior diameter of C1 (mm) (HIAPCI) (Figure 1).

3. Horizontal outer transverse diameter of C1 (mm) (HOTDCI) (Figure 1).
4. Distance between outer margin of transverse foramen and outer margin of lateral mass (mm) (outer margin) (Figure 1).
5. AP diameter of the superior surface of the C1 anterior arch (mm) (superior surface) as in Figure 1.
6. Lateral outer AP diameter of C1 (mm) (LOAPC1) (Figure 2).
7. Height of the atlas dorsal arch (mm) (dorsal arch) (Figure 2).
8. Frontal outer transverse diameter of C1 (mm) (FOTDCI) (Figure 3).
9. Angle along the axis line of the dens to the occlusal plane (dens angle (degrees) as in Figure 4).



Figure 1: Axial view of cervical vertebra.

(1) Horizontal outer anteroposterior (AP) diameter of the first cervical vertebra (C1) (mm) (HOAPCI). (2) Horizontal inner anteroposterior diameter of C1 (mm) (HIAPCI). (3) Horizontal outer transverse diameter of C1 (mm) (HOTDCI). (4) Distance between outer margin of transverse foramen and outer margin of lateral mass (mm) (outer margin). (5) AP diameter of superior surface of C1 anterior arch (mm) (superior surface).




Figure 2: Lateral view of second cervical vertebra.

No. (6)=the lateral outer AP diameter of C1 (mm) (LOAPC1). No. (7)=Height of the atlas dorsal arch (mm) (dorsal arch).

Ethical approval

This study was designed according to STROBE guidelines and conducted in accordance with the Declaration of Helsinki. The study protocol was approved by the Ethical Committee of the Faculty of Dentistry, Sana'a University, Yemen.

Statistical analysis

The statistical software tool SPSS (Statistical Tool for the Social Sciences) version 25 for Windows was used to enter and analyze the data. Means, standard deviations, chi-square and t-tests were calculated at 95% statistical significance level and a *p* value less than 0.05 was considered statistically significant.

RESULTS

The sample included 21 males and 31 females with no statistical significant difference between gender and both skeletal groups ($p>0.05$) (Table 1). Statistically significant differences were investigated between the first and third class groups. There was no statistically significant difference in the outer margin and dense angles ($p>0.05$). The variables HOAPC1, HIAPC1, HOTDC1, upper, LOAPC1, dorsal arch, and FOTDC1 showed statistically significant differences between the first class and the third class ($p<0.05$), and the highest mean value was found in the third class (Table 2).

DISCUSSION

Several previous studies have evaluated the maturity and shape of the cervical vertebrae using 3D imaging, but there has been insufficient research to evaluate the shape of the cervical vertebrae in Yemen and the Class I and III malocclusion patients. The majority of studies that used cone-beam CT were based on the evaluation of small samples due to the high cost despite the high accuracy of this method compared to 3D imaging^{1,20}. In this study, 52 patients were evaluated using cone-beam CT to determine the shape of the first cervical vertebrae in class I and III malocclusion in Yemen. This research study evaluated 52 patients with different skeletal patterns “class I and III,” which were analyzed based on gender. This was the same as that of Baydas *et al.*²¹, who evaluated the morphology of the cervical vertebrae of 90 patients (45 males and 45 females) with different skeletal classes. Furthermore, Gunduz *et al.*²²,

who examined the hyoid bone and C1 atlas morphology in patients with different skeletal classes, found that there was a statistically significant difference between genders.




Figure 3: Frontal view of second cervical vertebra.

No. (8)=the frontal outer transverse diameter of C1 (mm) (FOTDC1).




Figure 4: The angle along axis line of the dens to occlusal plane (dens angle [degrees]).

Table 1: The relation between gender and skeletal groups.

Categories	Gender		Total
	F	M	
Skeletal Class I	14	12	26
	53.8%	46.2%	100.0%
Class III	17	9	26
	65.4%	34.6%	100.0%
Total	31	21	52
	59.6%	40.4%	100.0%
p-value	0.719 (0.397)		

p: Probability; *: significance ≤ 0.05 ; Test used: Chi-Square test

Table 2: The mean value of cervical vertebra measurements in Class I and Class III groups.

Variables	Class I		Class III		t test	p-value	ICC	
	Mean	Std	Mean	Std			Class I	Class III
HOAPC1	37.35	3.35	40.08	2.57	-3.290	0.002**	0.95	0.96
HIAPC1	27.36	2.24	28.59	0.95	-2.557	0.015*	0.98	0.97
HOTDC1	68.00	3.19	72.53	3.40	-4.958	0.000**	0.94	0.95
Outer margin	13.33	1.76	13.36	1.74	-0.063	0.950	0.97	0.94
Superior surface	5.92	1.02	6.73	1.18	-2.672	0.010*	0.94	0.96
LOAPC1	37.99	2.95	40.14	2.63	-2.767	0.008**	0.93	0.94
DORSAL arch	7.74	1.38	8.65	1.04	-2.657	0.011*	0.90	0.91
FOTDC1	67.03	2.88	72.33	3.56	-6.384	0.000**	0.95	0.96
Dens Angle	-0.08	0.27	0	0	-1.443	0.161	0.98	0.97

HOAPC1: horizontal outer anteroposterior diameter of C1; HIAPC1: horizontal inner anteroposterior diameter of C1; HOTDC1: horizontal outer transverse diameter of C1; LOAPC1: lateral outer anteroposterior diameter of C1; FOTDC1: frontal outer transverse diameter of C1; SD: standard deviation, P: Probability; *: significance ≤ 0.05 ; Test used: Student's t-test unpaired

With the development of 3D imaging, it was found to be more reliable and accurate in detecting minor changes and evaluating cervical vertebrae than the

previously used 2D imaging techniques. This was confirmed by Patcas *et al.*²³, who suggested that CBCT images are needed for a more accurate diagnosis and

location of the deviations. In this study, CBCT images were evaluated for 52 patients from different skeletal classes to assess cervical vertebral morphology; this was the same as Watanabe *et al.*², and Ghazy *et al.*²⁴, who described the morphology of cervical vertebrae in different skeletal classes using CBCT. In this study, specific measurements were used to evaluate the morphology of the cervical vertebrae. The morphology of the cervical vertebrae was assessed using nine linear measurements and one angle assessment. It was discovered that patients from different skeletal classes did not differ statistically significantly in the outer margin and dense angles. This was in line with the findings of Ghazi *et al.*²⁴, who used the same methodology to examine the cervical vertebral morphology in patients with skeletal classes I and II, and Watanabe *et al.*², who examined the cervical vertebral morphology in patients with skeletal classes II and III. The variables in this study that demonstrated a statistically significant difference between Class I and Class III were HOAPC1, HIAPC1, HOTDC1, Superior, LOAPC 1, Dorsal arch, and FOTDC1. According to Watanabe *et al.*²², there were no appreciable changes between Class II and Class III patients' "HOTDC1," LOAPC1, FOTDC1, or "Dense Angle," but there was a notable difference between Class II and Class III patients' atlas "dorsal arch" heights. Given that the latter examined the morphology of the cervical vertebrae in Japanese patients, whereas this study focused on Yemeni patients, the disparity could be explained by the different heritages of the two groups. Additionally, this study contrasted Class III with Class I malocclusions, whereas their study contrasted Class III with Class II.

Limitations of the study

The limitations of this study include a small sample size because of CBCT cost and its limited use for orthodontic patients; Class II subjects were not included; current study focused on Class III and Class I because they are considered the most common among the Yemeni people, and Class II has been studied in previous research, and patients were not subdivided on their vertical skeletal patterns. For future studies, we recommend comparing the cervical vertebrae morphology in the different heritages all over the world.

CONCLUSIONS

This study confirmed previous findings that individuals with Class III and Class I malocclusions and different anteroposterior skeletal patterns had differences in cervical vertebral shape. No statistically significant difference was observed in the outer margin and dense angles, while HOAPC1, HIAPC1, HOTDC1, superior surface, LOAPC1, dorsal arch, and FOTDC1 showed statistically significant differences between individuals with different anteroposterior skeletal patterns. The measured parameters were found to be significantly increased in Class III patients compared to Class I.

ACKNOWLEDGEMENT

Thanks for Professor Elham Abualhija from Qatar University for her advices.

AUTHOR'S CONTRIBUTION

Al-Ghola E: writing original draft, methodology, investigation. **Al-Harazi G:** formal analysis, data curation, conceptualization, editing, methodology. Final manuscript was checked and approved by all authors.

DATA AVAILABILITY

The data that support the findings of this study are available from the corresponding author upon reasonable request.

CONFLICT OF INTEREST

None to declare.

REFERENCES

1. Manabe A, Ishida T, Kanda E, Ono T. Evaluation of maxillary and mandibular growth patterns with cephalometric analysis based on cervical vertebral maturation: A Japanese cross-sectional study. *Plos one* 2022; 17(4): e0265272. <https://doi.org/10.1371/journal.pone.0265272>
2. Watanabe M, Yamaguchi T, Maki K. Cervical vertebra morphology in different skeletal classes: A three-dimensional computed tomography evaluation. *The Angle Orthodontist* 2020; 80(4): 719-724. <https://doi.org/10.2319/100609-557.1>
3. Franchi L, Baccetti T, McNamara JA. Mandibular growth as related to cervical vertebral maturation and body height. *American J Orthod Dentofac Orthopedics* 2000; 118(3): 335-340. <https://doi.org/10.1067/mod.2000.107009>
4. Fransiska M, Retno W. Class III malocclusion camouflage treatment using a conventional orthodontic appliance in a non-growing patient. *J Dent Indonesia* 2023; 30(1):48-51. <https://doi.org/10.14693/jdi.v30i1.1467>
5. Martin CB, Chalmers EV, McIntyre GT, Cochrane H, Mossey PA. Orthodontic scanners: What's available? *J Orthodont* 2015; 42(2): 136-143. <https://doi.org/10.1179/1465313315Y.0000000001>
6. Baccetti T, Franchi L, McNamara Jr JA. The Cervical Vertebral Maturation (CVM) method for the assessment of optimal treatment timing in dentofacial orthopedics. *Seminars Orthodont* 2005;11(3): 119-29. <https://doi.org/10.1053/j.sodo.2005.04.005>
7. Lai E, Liu H, Chang JP, *et al.* Radiographic assessment of skeletal maturation stages for orthodontic patients: Hand-wrist bones or cervical vertebrae? *J Formosan Med Assoc* 2008; 107(4): 316-325. [https://doi.org/10.1016/S0929-6646\(08\)60093-5](https://doi.org/10.1016/S0929-6646(08)60093-5)
8. Zandi M, Shokri A, Mollabashi V, Eghdami Z, Amini P.. Anatomical characteristics of mandibular bone in skeletal class I, II and III patients by using cone beam computed tomography images in an Iranian population. *Brazilian Dental Sci* 2021; 24(2). <https://doi.org/10.14295/bds.2021.v24i2.2475>
9. Emre C, Orhan K, Misirli M, Bilecenoglu B. Cone beam computed tomography evaluation of the relationship between atlantodental interval and skeletal facial morphology in adolescents. *Brazilian J Otorhinolaryngol* 2020; 86(6): 711-719.

- <https://doi.org/10.1016/j.bjorl.2019.05.005>
10. Abdul-Majid ALA, Aleryani HAA, Aldeen YAAS, Al-Shamahy HA. Reliability of modern radiographic dental age estimation methods among Yemeni children in Sana'a city, Yemen. *Universal J Pharm Res* 2025; 9(6): 14-24. <https://doi.org/10.22270/ujpr.v9i6.1234>
 11. Al-Subbary IA, Obeyah AA, Al-Mogahed NM, Al-Ammari MH. Dental caries and treatment needs among children with physical disabilities in Dhamar city, Yemen: A comparative study. *Universal J Pharm Res* 2024; 9(2):34-39. <https://doi.org/10.22270/ujpr.v9i2.1086>
 12. Sharafuddin AH, Alshameri BH, AL-Haddad KA, Al-Najhi MMA, Al-Shamahy HA. The effect of dental implants on increasing the colonization rate of aerobic bacteria in the oral cavity. *Universal J Pharm Res* 2023; 8(3): 28-33. <https://doi.org/10.22270/ujpr.v8i3.944>
 13. Al-Hamzi MA, Sharafuddin AH, Al-Shamahy HA, *et al.* The effect of dental implants on aerobic bacteria colonization in the oral cavity and the antibiotic profile of common isolated aerobic bacteria. *Universal J Pharm Res* 2023; 8(4): 1-8. <https://doi.org/10.22270/ujpr.v8i4.969>
 14. Sharaf-Aldeen HMA, Abbas AKMA, Al-Shamahy HA, *et al.* Oromandibular dystonia: Prevalence, clinical and demographic data, therapeutic strategies out-come for hundred patients in Sana'a city, Yemen. *Universal J Pharm Res* 2023; 8(2): 61-70. <https://doi.org/10.22270/ujpr.v8i2.925>
 15. Alsada'a WAA, Al-Hamzi MA, Al-Khawlani AI, Al-Shamahy HA. Radiographic evaluation of overhanging restorations among Yemeni patients using cone-beam computed tomography (CBCT). *Universal J Pharm Res* 2023; 8(5): 30-38. <https://doi.org/10.22270/ujpr.v8i5.1007>
 16. Zabara AQMQ, Al-Kholani AIM, Al-Shamahy HA, *et al.* Resolution of factors and pattern of permanent dental extraction in selected dental clinics in Sana'a city, Yemen. *Universal J Pharm Res* 2022; 7(4): 44-49. <https://doi.org/10.22270/ujpr.v7i4.813>
 17. Al-Khorasani MAM, Al-Kebsi AM, Al-Shamahy HA, *et al.* Prevalence of signs of temporomandibular disorders in healthy asymptomatic completely edentulous individuals and the effect of denture on temporomandibular disorders. *Universal J Pharm Res* 2023; 8(1):28-34. <https://doi.org/10.22270/ujpr.v8i1.894>
 18. Dahag WAM, Al-Kholani AIM, Al-Shamahy HA, *et al.* Tanaka and Johnston's mixed dentition validity: An analysis among Yemeni adults in Sana'a city. *Universal J Pharm Res* 2022; 6(6): 1-5. <https://doi.org/10.22270/ujpr.v6i6.691>
 19. Daer AA, Abuaffan AH. Cephalometric norms among a sample of Yemeni adults. *Orthodontic Waves* 2016;75(2):35-40. <https://doi.org/10.1016/j.odw.2016.03.00>
 20. Sonnesen L, Kjaer I. Cervical vertebral body fusions in patients with skeletal deep bite. *The European J Orthodont* 2007; 29(5): 464-470. <https://doi.org/10.1093/ejo/cjm043>
 21. Baydaş B, Yavuz I, Durna N, Ceylan I. An investigation of cervicovertebral morphology in different sagittal skeletal growth patterns. *The European J Orthodont* 2004; 26(1): 43-49. <https://doi.org/10.1093/ejo/26.1.43>
 22. Gündüz AS, Dildeş N, Devcioglu KJ. Cephalometric investigation of first cervical vertebrae morphology and hyoid position in young adults with different sagittal skeletal patterns. *The Sci World J* 2014; 159784. <https://doi.org/10.1155/2014/159784>
 23. Patcas R, Tausch D, Pandis N, *et al.* Illusions of fusions: Assessing cervical vertebral fusion on lateral cephalograms, multidetector computed tomographs, and cone-beam computed tomographs. *American J Orthodont Dentofacial Orth* 2013; 143(2), 213-220. <https://doi.org/10.1016/j.ajodo.2012.09.017>
 24. Ghazy A, Refaat WE, Morcos SS. Three-dimensional evaluation of cervical vertebral morphology in skeletal class II of malocclusion in Egyptians. *Suez Canal Univ Med J* 2017; 20(1): 62-67. <https://doi.org/10.21608/scumj.2017.46988>